INSTRUCTIONS FOR EPIDURAL STERIOD INJECTION (ESI)

- 1. Your procedure is scheduled for ______ Arrival time ______ Please report to Orthopedic Surgery Center on the first floor.
- 2. **Nothing to eat or drink** after midnight the night before your procedure. If you need to take regularly scheduled medications only a sip of water is allowed.
- 3. You must have a driver present with you.
- 4. The following medications must be discontinued before an ESI can be performed:

All Non Steriodal Anti-Inflammatories (NSAIDS) must be stopped 2 days before injection:

Advil (Ibuprofen, motrin)	Feldene (piroxicam)	
Aleve (naproxen)	Daypro (oxaprozin)	
Mobic (meloxicam)	Athrotec (diclofenac +misoprostol)	
Voltaren (diclofenac)	· ,	
Relafen (nabumetone)	Indocin (Indomethacin)	
Lodine (etodolac)	Orudis (ketoprofen)	
× ,	Toradol (ketorolac)	

Clinoril (sulindac)

All Asprin containing products must be stopped 5 days before injection:

Asprin 81 mg	Fiorinal
Excedrin	Bufferin
Asprin 325 mg	Goody's powder

All blood thinners must be stopped 7 days before injection:

Coumadin (warfarin) Persantine (dipyridamole) Plavix (clopidogrel) Aggrenox (asprin +

dipyridamole)

Lovenox (enoxarain) must be stopped 2 days before injection

Please avoid all herbal medications including vitamin E and fish oil 2 days before injection. You may resume your medications 2 days after your procedure date.

- 5. If you are on aspirin, coumadin or plavix, specific blood work will need to be obtained before your injection.
- 6. If you are older than 50 years of age or have a cardiac history, we must have an EKG within the last 6 months.
- 7. If a medical or cardiac clearance is needed, we must have the clearance before your procedure is scheduled.
- 8. The ESI may take 2-7 days to take full effect. You may still continue to have pain immediately following the injection, this is normal.
- 9. Please call our office (833-774-6352) if you experience severe pain, fever >100.4 or a severe headache associated with nausea and vomiting.
- 10. If you do not hear from Orthopedic Surgery Center at least one day before your injection contact Dr. McCarthy's office (833) 774-6352 Ext 21

Please sign below acknowledging you understand the above instructions and directions and have no questions.

X ----- Date